Screening of Colorectal Cancer: Progress and Problems

Sidney J. Winawer

Memorial Sloan-Kettering Cancer Center, 1275 York Ave., New York, NY 10021, USA

1	Introduction	231
2	Progress.	233
2.1	Guidelines Consensus	233
2.2	Fecal Occult Blood Testing	233
2.3	Sigmoidoscopy	235
2.4	Screening Colonoscopy	236
2.5	Adenomas.	236
3	Problems	238
3.1	Screening Rates.	238
3.2	Implementation	238
3.3	Barriers	240
4	Concluding Remarks	241
	References	242

Abstract Considerable progress has been made in the past three decades in our understanding of the biology and prevention of colorectal cancer. The long natural history of colorectal cancer as it evolves from adenomatous polyps in the majority of cases provides opportunities for detection of early stage cancer and for prevention of cancer by removal of adenomas. Strong evidence of the effectiveness of screening has resulted in a world-wide consensus, as reported in evidence-based guidelines, that screening should be offered to all men and women age 50 and older, younger in the presence of factors that increase risk. Several options are now available for screening, and the emerging technology of stool DNA testing and virtual colonoscopy shows promise. However, many problems remain to be addressed. Screening rates are low. Successful strategies need to be implemented to overcome patient and system barriers. Resources, especially endoscopic capacity, may be inadequate to handle the burden of screening, diagnosis, and follow-up surveillances. There are quality-control issues at every step. Stratification of people by risk, a two-stage screening approach and less intensive surveillance following polypectomy can be helpful. Colorectal cancer screening is cost-effective and could save many lives each year if it were widely implemented.

1 Introduction

Colorectal cancer accounts for approximately 945,000 new cases and 500,000 deaths each year worldwide [1] (to put those numbers in context, see

Table 1. The burden of cancer (both sexes) in the more developed countries. Estimates in 2000 and expected cases in 2020 [1]

	Estimates in 2000			Expected cases in 2020 (<i>n</i>)
	Cases (<i>n</i>)	Deaths (<i>n</i>)	Ratio of cases:deaths	
Large bowel	610,591	301,648	2.02	806,176
Stomach	334,011	229,939	1.45	439,842
Pancreas	127,416	128,730	0.98	168,453
Liver	106,950	105,649	1.01	142,836
Esophagus	71,163	63,938	1.11	94,251
Total	1,250,131	829,904	1.50	1,651,558

Table 2. The burden of cancer (both sexes) in less developed countries. Estimates in 2000 and expected cases in 2020 [1]

	Estimates in 2000			Expected cases in 2020 (<i>n</i>)
	Cases (<i>n</i>)	Deaths (<i>n</i>)	Ratio of cases:deaths	
Stomach	543,026	416,614	1.30	983,414
Liver	457,406	442,916	1.03	801,685
Esophagus	341,163	273,560	1.24	624,574
Large bowel	334,123	190,761	1.75	162,401
Pancreas	88,969	84,734	1.05	162,401
Total	1,764,687	1,408,585	1.25	3,164,220

Tables 1 and 2). Considerable progress has been made in the past 25 years in our ability to reduce this burden, with the introduction of technology for screening, early diagnosis, and prevention by removal of the premalignant polyp [2]. During this period of time, greater insight into the adenoma–carcinoma sequence has provided us with an understanding of the progress made and the problems of screening. The risk for colorectal cancer in average-risk men and women, as well as in groups at increased risk, has also been clarified. Strong evidence has accumulated that screening is effective in reducing the incidence and mortality of colorectal cancer. Although tremendous progress has been made over the last 25 years in these areas, we now face many problems, including the challenging prospects of universal implementation of effective screening methods and integration of new technology and concepts as they develop [2].

2 Progress

2.1 Guidelines Consensus

There was considerable controversy as to whether screening was effective, but beginning in 1996 a consensus evolved reflected in positive guidelines by a number of evidence-based assessments by authoritative bodies including the U.S. Preventative Services Task Force, the Agency for Health Care Policy and Research, the American Cancer Society, World Health Organization, Ontario Expert Panel, Australian Task Force, and the European Screening Group, just to cite a few [2]. The U.S. Preventive Services Task Force upgraded its recommendation to Grade A [3]. The consensus was that all men and women should be screened for colorectal cancer and adenomatous polyps beginning at age 50, or younger in the presence of factors that increase risk.

The majority of patients who will develop colorectal cancer are considered to be at average risk (75%) with no special risk factors. People at increased risk include those with inflammatory bowel disease, which is associated with 1% of the colorectal cancers each year; familial adenomatous polyposis (FAP), 1%; hereditary non-polyposis colorectal cancer (HNPCC), 2%–5%; and those who have a family history consisting of one or two close relatives, 15%–20% [4]. We have a much better understanding today of the magnitude and clinical patterns in these increased risk groups. People having one first degree relative with either a colorectal cancer or an adenomatous polyp at a young age have approximately a twofold increased risk of colorectal cancer, and the risk begins about 10 years younger than in average-risk patients. The risk in close relatives increases inversely with the age in which the proband had an adenomatous polyp or colorectal cancer, the younger the age of the proband the greater the risk in the first-degree relatives [5].

2.2 Fecal Occult Blood Testing

The early evidence of the benefit of colorectal cancer screening with fecal occult blood testing demonstrated a shift in screen detected cancers to an earlier stage with fewer Dukes' D screen-detected cancers as compared to controls [6]. However, this stage shift was quickly challenged as possibly due to length bias and lead time bias. Length bias suggests that screening would pick up cancers that are slow growing more often than cancers that are more aggressive, and lead time bias suggests that screening merely detects the cancers earlier in their natural history, but with no change in their outcome [4]. However, more recently, results of three major randomized control trials

Table 3. Fecal occult blood test (FOBT) screening in randomized controlled trials [7–9]

Trial	Mortality reduction (%)		
	Biennial	Annual	Compliers
Minnesota ^a 47,000/18 years	21	33	45
Denmark 140,000/10 years	18	-	30
UK 153,000/7.8 years	15	-	-

^a Sensitive slide test used.

demonstrated a mortality reduction in colorectal cancer of the entire screened cohort (Table 3). The largest reduction in mortality was in the University of Minnesota program, in which one group of patients was screened annually with a sensitive slide test. That group demonstrated a 33% reduction in mortality [7]. The compliers with the test demonstrated a 45% reduction in mortality.

The Minnesota Colon Cancer Control Study is a long-term, prospective, randomized, controlled trial of stool blood test efficacy in the detection of colorectal cancer [7]. In this trial, 46,551 participants between 50 and 80 years of age were assigned to either a study group that was offered annual stool blood testing, a second study group that was offered biennial stool blood testing, or a third control group that was not offered screening. The majority (approximately 83%) of stool slides used were rehydrated. The compliance rate for stool slide preparation was approximately 75%. The overall test positivity rate was initially 2.4% with nonrehydrated slides, and it subsequently increased to 9.8% with rehydrated slides. Nonrehydrated slide-test programmatic sensitivity was 80% and specificity was 98%, whereas rehydrated slide-test sensitivity increased to 92% and specificity decreased to 90%. The test's positive predictive value for colorectal adenomas and cancer was 31%. Patients with positive stool blood tests had a diagnostic evaluation with colonoscopy. During a 13-year follow-up period, this trial demonstrated that annual stool blood testing resulted in a significant 33% reduction in colorectal cancer mortality, an improved survival in those individuals diagnosed with colorectal cancer, and a shift towards detection of earlier-stage cancers. A recent update of this trial through year 18 demonstrated a 36% reduction in colorectal cancer mortality with annual stool blood testing [7].

A large United Kingdom population-based, randomized, controlled trial of screening with stool blood testing was performed in asymptomatic individuals between 50 and 74 years of age who were selected from general practitioner lists in the Nottingham area of England [8]. Of the 156,000 participants recruited, 52,258 individuals were randomized to the study group who were offered biennial screening with nonrehydrated stool blood testing. The initial compliance rate for stool slide preparation was 53%. The overall test

positivity rate was 2.3%. The slide-test sensitivity was 72% and specificity was 98%. The test's positive predictive value for colorectal adenomas and cancer was 53%. Patients with positive stool blood tests had a diagnostic evaluation with colonoscopy. An increased number of earlier-stage (Dukes' A or B) cancers were detected in the study group (90% versus 40%) as compared to the control group. The trial demonstrated a 15% reduction in colorectal cancer mortality, with a median follow-up of 7.8 years (odds ratio, 0.85) [8].

A large Danish population-based, randomized, controlled trial of stool blood test screening randomized 62,000 participants between 45 and 74 years of age to either biennial nonhydrated stool testing or a control group [9]. The initial compliance rate for stool slide preparation was 67%. The overall test positivity rate was 1.0%. Patients with positive stool blood tests had a diagnostic evaluation with colonoscopy. This trial, now with 10 years of follow-up data, reported an 18% reduction in colorectal cancer mortality, including deaths from complications related to colorectal cancer treatment, in the screened group (mortality ratio, 0.82). All of the trials in the United States and Europe are consistent with one another in terms of the magnitude of the mortality reduction as a function of the type of slide test used and the frequency of screening. This put to rest questions of screening bias, and for the first time convincingly demonstrated that screening for colorectal cancer with fecal occult blood testing is effective.

2.3

Sigmoidoscopy

Colorectal cancer mortality was also shown to be reduced by sigmoidoscopy in two-case control studies. A retrospective, case-control study from the Kaiser Permanente group has provided strong evidence that sigmoidoscopy can significantly reduce rectosigmoid cancer mortality [10]. This study compared the use of screening rigid sigmoidoscopy during the 10-year period prior to diagnosis in 261 individuals who died of rectosigmoid cancer with 868 matched controls. The results showed that only 8.8% of case patients had a screening sigmoidoscopy, as compared to 24.2% of the controls, with an adjusted odds ratio of 0.41, and that this negative association remained as strong with screening intervals from prior sigmoidoscopy to cancer diagnosis of as long as 9–10 years. This study demonstrated a 60% reduction in rectosigmoid cancer mortality.

A second smaller case-control study of screening sigmoidoscopy from the University of Wisconsin [11], which compared the records of 66 individuals of the Greater Marshfield Community Health Plan who died of colorectal cancer with 196 matched controls, found that case patients were less likely to have had a screening sigmoidoscopy than were controls (10% versus

30%). This study also demonstrated that a single screening flexible sigmoidoscopy resulted in a reduction in colorectal cancer mortality.

In 1993 the National Cancer Institute initiated a large, multicenter, long-term, randomized, controlled screening trial (PLCO trial: prostate, lung, colorectal, ovary) of 148,000 individuals that includes flexible sigmoidoscopy (60-cm sigmoidoscope) for colorectal cancer screening [12]. A second large, multicenter clinical trial to assess the efficacy of screening flexible sigmoidoscopy has been organized in England [13]. No long-term trials utilizing flexible sigmoidoscopy have yet reported incidence or mortality data. The combination of fecal occult blood testing added to sigmoidoscopy has also been associated with a mortality reduction in a study conducted in the U.S. [14]. As a result of these studies, colorectal cancer screening guidelines recommend that average-risk men and women age 50 or older (or age ≥ 40 with a family history) be offered options for screening which include either fecal occult blood testing annually or flexible sigmoidoscopy every 5 years, or a combination of the two [2, 3, 4].

2.4

Screening Colonoscopy

Although no studies in average-risk individuals have evaluated whether screening colonoscopy alone can reduce colorectal cancer mortality or incidence, some evidence suggests that this examination should be considered for colorectal cancer screening in this population. In comparison to fecal occult blood testing, which has a low sensitivity for the detection of adenomas, and to sigmoidoscopy, which examines at most only the distal third of the large bowel, colonoscopy (1) is very sensitive for the detection of cancer and both small and large adenomas, (2) completely examines the entire colon and rectum, and (3) provides the opportunity for the endoscopic removal of adenomas and biopsy of suspicious mass lesions.

There have been two major studies that have reported results of screening colonoscopy, one a U.S. Veterans Affairs Cooperative study [15], and another conducted in employees of Eli-Lilly. The total number of patients examined in both studies was more than 5,000, and the studies were similar in outcome, demonstrating that about 10% of asymptomatic patients over the age of 50, both men and women, have either a cancer or an advanced adenoma [15, 16].

2.5

Adenomas

The most frequent outcome of screening is not cancer, but an adenomatous polyp [17]. These polyps can be removed quite expeditiously by colonoscopy as an outpatient procedure—with the entire colon being examined—in

15–20 min. This was a major advance compared to the exploratory laparotomy and multiple colostomies that were done in the past. The feasibility of removing polyps through the colonoscope was reported in the mid-1970s [18], and shortly thereafter the National Polyp Study was organized as a randomized, multicenter, controlled trial to examine surveillance intervals and methods after polypectomy, to examine the potential incidence reduction of colorectal cancer following polypectomy, and to study the adenoma–carcinoma natural history and biology [19]. The National Polyp Study demonstrated that although many polyps were found at follow-up colonoscopy, the initial colonoscopy was successful in clearing out the colon of advanced adenomas [20]. Advanced adenomas are defined as those 1 cm in size or larger or with high-grade dysplasia or invasive cancer, and some investigators also include in the definition a high degree of villous component [21]. In the National Polyp Study, it was demonstrated that at a 3-year follow-up examination, only 3% of the patients had advanced adenomas after they had their colons cleared of all polyps at baseline. An examination at 1 year before the 3-year examination provided no additional benefit. This resulted in guidelines that now recommend omitting the 1-year examination after polypectomy and going to a 3-year examination for the first follow-up [2, 3, 4].

It was long believed that the adenoma was the precursor of colorectal cancer and its removal would result in the prevention of colorectal cancer [18]. This belief was substantiated by data from the National Polyp Study, which demonstrated a reduction (70%–90%) in observed cancers as compared to expected cancers in the National Polyp Study cohort following the baseline clearing colonoscopy [22]. This was later confirmed in an Italian multicenter study (66% incidence reduction) [23] (Table 4). The prevention of colorectal cancer by polypectomy is one of the best-kept secrets from the public. It is one of the most powerful prevention strategies that is available, and yet it is not widely known. In the National Polyp Study, it was further demonstrated that one could stratify patients into those at high risk for advanced adenomas in the future as compared to those at low risk for advanced adenomas. The low risk patients constituted 70% of the cohort and could have their first follow-up colonoscopy years after their clearing colonoscopy rather than at 3 years. It is reasonable to do the first follow-up colonoscopy in these patients at 5 years and reserve the three-year follow-up colonoscopy for only high-risk patients. This high-risk group is defined as those who have multiple adenomas at baseline or have a positive family history and are at age

Table 4. Effect of colonoscopic polypectomy on incidence of colorectal cancer [22, 23]

Study	↓ Incidence (%)
U.S. National Polyp Study	76–90
Italian Multicenter Study Group	66

over 60 at their first polypectomy. This stratification would reduce complications and costs, and conserve resources, which could be better directed towards the initial screening of patients [2].

3 Problems

3.1 Screening Rates

Screening rates for colorectal cancer are low and well below those for mammography. National Health Interview Surveys have consistently demonstrated this fact. In the U.S., only 20% of people over age 50 have had fecal occult blood test, and one out of three has had an endoscopic screening test in the previous 5 years [24]. One of the most frequent reasons for not getting screened (88%), is that “it was not recommended by my doctor,” according to one survey [18]. The health care provider plays a very important role in motivating patients to be screened. Health care providers may be more motivated today because of the litigation that is now occurring in colorectal cancer with the most frequent reasons for litigation being failure to screen, failure to diagnose, and deviation from standard of care. Standard of care now is based on published guidelines rather than practice in the community. These goals have been enhanced recently by U.S. Congress legislation, which in the year 2000 provided Medicare reimbursement for fecal occult blood testing, flexible sigmoidoscopy, and, beginning in July 2001, Medicare reimbursement for screening colonoscopy. This legislation was based on the demonstration in recent years by many cost-effectiveness models, that colorectal cancer screening is cost-effective; costing less than US \$20,000 per life-year saved, and is equivalent to screening mammography. Colorectal cancer screening has further been enhanced by the designation of March as National Colorectal Cancer Awareness Month, by former U.S. President Bill Clinton [18].

3.2 Implementation

Universal implementation of screening is a major challenge. In the U.S. the target population is 70 million men and women age 50 and over. The question is whether there are resources to accomplish screening in such a large population. There are several alternatives that can be considered. Direct population screening colonoscopy is one strategy. We can identify those people who have adenomas, who can then be further stratified into low and

high risk for subsequent advanced adenomas, with less intense or more intense follow-up surveillance, respectively.

Another approach is to try to identify those individuals who would be most likely to benefit from screening. One such approach is through genetic testing. However, genetic testing can identify only 5%–6% of those people destined to get colorectal cancer, and these individuals must first be identified by their family history of either FAP or HNPCC [25]. New genetic mutations in colorectal cancer have been demonstrated in the general population; two in Ashkenazi Jews and one in non-Jewish people. The genetic mutations in Ashkenazi Jews have a very low penetrance, and by themselves do not confer a striking increase in risk. They probably require additional polymorphisms to substantially increase the risk for colorectal cancer in those individuals [25]. There has been recent interest in studying genetic mutations in the stool, recent reports demonstrating that a panel of DNA mutations in stool could result in the detection of cancers and adenomas with a sensitivity of 50%–60% and specificity of over 90% [26–29]. The best stool marker that we have right now is the fecal occult blood test, which can reduce mortality if done on a regular basis annually. This can be done with either the guaiac-based tests or by an immunochemical test (Table 5) [30, 31]. In the two-stage strategy, screening sigmoidoscopy, a stool marker, or virtual colonoscopy would be used first. Only those individuals having a positive significant lesion found would be referred for fiberoptic colonoscopy with biopsy or polypectomy. The accuracy of virtual colonoscopy has not yet been well established, and it is not totally non-invasive [32, 33]. The sensitivity has ranged from 1/3 to over 90% for polyps over 1 cm in size. Patients require a preparation for 1 day before the examination, and have air instilled in the bowel. This is, however, promising and needs further study. There have been many advances recently in diagnostic endoscopy with magnification lenses, dye staining, spectroscopy, and optical coherence tomography, which would greatly enhance our ability to make tissue diagnosis more accurately and perhaps separate those patients endoscopically who do not need biopsy or polypectomy, especially in the case of very small polyps [18].

Table 5. Choice of fecal occult blood test (FOBT) [30]

Colonoscopy resources	Population compliance with diet and drug restrictions for guaiac test	Suggested FOBT
Limited	Reliable	High specificity guaiac test (e.g., Hemoccult)
Limited	Unreliable or uncertain	Immunochemical test
Readily available	Reliable	Sensitive guaiac test (e.g., Hemoccult Sensa)
Readily available	Unreliable or uncertain	Immunochemical test

The resource issue is critical. Endoscopy centers are being overwhelmed by the number of referred colonoscopies. These include screening colonoscopy, diagnostic colonoscopies in patients with a positive screening test, and colonoscopy in symptomatic patients. Endoscopic resources could be freed up for screening and diagnosis if the post-polypectomy surveillance interval could have reduced intensity. Patients can be stratified after initial polypectomy into those who are at low or high risk for advanced adenomas later. Those at low risk could have their first post-polypectomy examination deferred for 5 years rather than 3 years. Guidelines have been modified recently to make this change in recommendations [2].

3.3

Barriers

Many barriers exist that prevent widespread implementation of screening, including financial concerns, patient inconvenience, and patient perceptions of benefit, risk, and discomfort [34, 35]. For screening programs to be successful, a cascade of events must be negotiated from beginning to end [2]. Physicians must remember to offer screening, patients must accept this advice, insurers must pay for screening and follow-up testing, and patient care organizations must have systems to track whether screening has taken place and provide reminders if it has not. Screening examinations must be feasible for providers, which is a special problem for sigmoidoscopy, and the workforce to do examinations well must be in place, a problem for colonoscopy. If any one stage in this sequence is faulty, the screening program will fail. Therefore, those who care about effective screening programs must be concerned with all of these elements of success.

The number of places where breakdowns can occur is large. Some patients may not understand or carry out bowel preparation instructions. Providers must be able to perform tests correctly. Office staff, aided by information systems, must remember when screening tests are due and patients must accept part of this responsibility because they commonly change providers (because their health plan changes) or move out of the area, leaving new doctors unable to determine when 5–10 years have passed since the last endoscopy. Also, shared decision-making can be difficult to implement. Not all patients want to share decisions and many prefer doctors to make a recommendation. Physicians may lack time, skill, and resources to carry out shared decision-making correctly, and patients may not be able to digest the information and information presented. Many forms of patient information about colorectal cancer are available [2].

4

Concluding Remarks

Cost-effectiveness analyses have shown that the cost per year of life saved by screening with any of the tests recommended is reasonable by U.S. standards [2, 36–40]. Although the specific results vary among analyses, in general the marginal cost-effectiveness of this screening is less than US \$25,000 per year of life saved. Screening for colorectal cancer was among the highest ranked services in an analysis of the value of preventive services based on the burden of disease prevented and cost effectiveness.

Though the up-front costs vary by screening modality, the long-term cost-effectiveness is similar across screening tests, so that decisions about which options to include, in the long run and from the perspective of society, do not need to be heavily affected by costs. Costs increase out of proportion to benefits with shorter intervals between screening examinations.

Screening has provided great opportunities. Screening can prevent colorectal cancer by polypectomy and find early-stage cancers for treatment with less morbidity. Screening can reduce the burden of treating advanced cancers, and can identify families at increased risk. Screening has also provided a better understanding of the biology of colorectal cancer [41].

Screening for colorectal cancer should be part of a complete prevention program, which includes a healthy lifestyle and familial risk assessment. Those individuals with increased familial risk require special screening approaches while those at average risk can have more standard screening. The average-risk individuals can be further stratified into those that require intensive follow-up and those who require less intensive or no follow-up at all. We are beginning to learn how to apply screening and surveillance approaches based on risk stratification for a more cost-effective approach in order to conserve resources and reduce complications and costs. Chemoprevention can be added to the program when substantial benefit of agents has been demonstrated. We now have a better understanding of the biology of colorectal cancer and the technology to intervene in that biology to make a difference in the lives of many people. We have the concepts and technology today to substantially reduce the mortality for colorectal cancer and even entirely prevent it [41].

Newer screening tests, or others yet to be developed, may with time replace the present options. Nevertheless, screening should take place with the tests available now and not wait until something better comes along. In this way, needless suffering and loss of life can be avoided for this leading cause of cancer death. Screening may become even more successful if the promise of new technologies is confirmed and they enter clinical practice. In the last analysis, the best test is the one that gets done, and gets done now.

References

1. René Lambert, in collaboration with the IDC (2003) An overview of the epidemiology and prevention of digestive cancer. *World Gastroenterol News* 8:21–25. Available at http://www.omge.org/publications/archive/2003_2/idca/idca2.htm. Cited 11 June 2004
2. Winawer SJ, Fletcher R, Rex D, Bond J, Burt R, et al (2003) Colorectal cancer screening and surveillance: clinical guidelines and rationale—update based on new evidence. *Gastroenterology* 124:544–560
3. US Preventive Services Task Force (2002) Screening for colorectal cancer: recommendations and rationale. *Ann Intern Med* 137:129–131
4. Winawer SJ, Fletcher RH, Miller L, Godlee F, Stolar MH, Mulrow CD, et al (1997) Colorectal cancer screening: clinical guidelines and rationale. *Gastroenterology* 112:594–642
5. St John JB, McDermott FT, Hopper JL, Debney EA, Johnson WR, Huges ES (1993) Cancer risk in relatives of patients with common colorectal cancer. *Ann Intern Med* 118:785–790
6. Kewenter J, Brevinge H, Engaras B, et al (1994) Results of screening, rescreening, and follow-up in a prospective randomized study for detection of colorectal cancer by fecal occult blood testing. Results for 68,308 subjects. *Scand J Gastroenterol* 29:468–473
7. Mandel JS, Bond JH, Church TR, Snover DC, et al (1993) Reducing mortality from colorectal cancer by screening for fecal occult blood. Minnesota Colon Cancer Control Study. *N Engl J Med* 328:1365–1371
8. Hardcastle JD, Chamberlain JO, Robinson MH, Moss SM, Amar SS, Balfour TW, et al (1996) Randomised controlled trial of faecal-occult-blood screening for colorectal cancer. *Lancet* 348:1472–1477
9. Kronborg O, Fenger c, Olsen J, et al (1996) Randomised study of screening for colorectal cancer with faecal-occult-blood test. *Lancet* 348:1467–1471
10. Selby JV, Friedman GD, Quesenberry CP Jr, Weiss NS (1992) A case control study of screening sigmoidoscopy and mortality from colorectal cancer. *N Engl J Med* 326:653–657
11. Newcomb PA, Norfleet RG, Storer BE, Surawicz TS, Marcus PM (1992) Screening sigmoidoscopy and colorectal cancer mortality. *J Natl Cancer Inst* 84:1572–1575
12. Gohagen JK, Prorok PC, Kramer BS, et al (1995) The prostate, lung, colorectal, and ovarian cancer screening trial of the National Cancer Institute. *Cancer* 75:1869–1873
13. Atkin WS, Edwards R, Wardle J, Northover JM, Sutton S, Hart AR, Williams CB, Cuzick J (2002) Single flexible sigmoidoscopy screening to prevent colorectal cancer: baseline findings of a UK multicentre randomized trial. *Lancet* 359:1291–1300
14. Winawer SJ, Flehinger BJ, Schottenfeld D, Miller DG (1993) Screening for colorectal cancer with fecal occult blood testing and sigmoidoscopy. *J Natl Cancer Inst* 85:1311–1318
15. Lieberman DA, Weiss DG, Bond JH, et al (2000) Use of colonoscopy to screen asymptomatic adults for colorectal cancer. Veterans Affairs Cooperative Study Group 380. *N Engl J Med* 343:162–168
16. Imperiale TF, Wagner DR, Lin CY, Larkin GN, Rogge JD, Ransohoff DF (2000) Risk of advanced proximal neoplasms in asymptomatic adults according to the distal colorectal findings. *N Engl J Med* 343:169–174
17. Winawer SJ, Fletcher RH, Miller L, et al (1997) Colorectal cancer screening: clinical guidelines and rationale. *Gastroenterology* 112:594–642
18. Winawer SJ (2001) American Cancer Society Award Lecture. A quarter century of colorectal cancer screening: progress and prospects. *J Clin Oncol* 19:6s–12s

19. O'Brien MJ, Winawer SJ, Zauber AG, et al (1990) The National Polyp Study: patient and polyp characteristics associated with high-grade dysplasia in colorectal adenomas. *Gastroenterology* 98:371–379
20. Winawer SJ, Zauber AG, O'Brien MJ, Nah MN, et al (1993) Randomized comparison of surveillance intervals after colonoscopic removal of newly diagnosed adenomatous polyps. *N Engl J Med* 328:901–906
21. Winawer SJ, Zauber AG (2002) The advanced adenomas as the primary target of screening. *Gastrointest Endosc Clin N Am* 12:1–9
22. Winawer SJ, Zauber AG, H MN, et al (1993) Prevention of colorectal cancer by colonoscopic polypectomy. *N Engl J Med* 329:1977–1981
23. Citarda F, Tomaselli G, Capocaccia R, Barcherini S, Crespi M (2001) The Italian multicenter study group. Efficacy in standard clinical practice of colonoscopic polypectomy in reducing colorectal cancer incidence. *Gut* 48:812–815
24. [No authors listed] (2001) Trends in screening for colorectal cancer—United States, 1997 and 1999. *MMWR Morb Mortal Wkly Rep* 50:162–166
25. Burt RW (2000) Colon cancer screening. *Gastroenterology* 119:837–853
26. Ahlquist DA, Skoletsky JE, Boynton KA, Harrington JJ, Mahoney DW, Pierceall WE, Thibodeau SN, Shuber Ap (2000) Colorectal cancer screening by detection of altered human DNA in stool: Feasibility of a multitarget assay panel. *Gastroenterology* 119:1219–1227
27. Dong SM, Traverso G, Johnson C, Geng L, Favis R, Boynton K, et al (2001) Detecting colorectal cancer in stool with the use of multiple genetic targets. *J Natl Cancer Inst* 93:858–865
28. Traverso G, Shuber MS, Levin B, Johnson, C, et al (2002) Detection of APC mutations in fecal DNA from patients with colorectal tumors. *N Engl J Med* 346:311–320
29. Traverso, G, Suber A, Olsson L, Levin B, Johnson C, et al (2002) Detection of proximal colorectal cancer through analysis of faecal DNA. *Lancet* 359:403–404
30. Young GP, St John DJ, Winawer SJ, Rozen P; WHO (World Health Organization) and OMED (World Organization for Digestive Endoscopy) (2002) Choice of fecal occult blood tests for colorectal cancer screening: recommendations based on performance characteristics in population studies. A WHO (World Health Organization) and OMED (World Organization for Digestive Endoscopy) report. *Am J Gastroenterol* 97:2499–2507
31. Rozen P, Knaani J, Samuel Z (2000) Comparative screening with a sensitive guaiac and specific immunochemical occult blood test in an endoscopic study. *Cancer* 89:46–52
32. Rex D (2002) Considering virtual colonoscopy. *Rev Gastroenterol Disord* 2:97–105
33. Pickhardt P, Choi R, Hwang I, Butler J, et al (2003) Computed tomographic virtual colonoscopy to screen for colorectal neoplasia in asymptomatic adults. *N Engl J Med* 349:2191–2200
34. Carlson MDA, Zauber AG, Winawer SJ, Crespi M, Peter C, Rosen P (2001) Lack of financial coverage is a major barrier to colorectal cancer screening: international survey results. *Am J Epidemiol* 153:s249 (abstract 926)
35. Winawer SJ, Zauber AG, Carlson MDA, Crespi M, Rozen P (2001) OMED/WHO/ASGE Worldwide Project for the Prevention of Colorectal Cancer: results of an international survey (abstract). *Gastrointest Endosc* 53:AB188
36. Wagner JL, Tunis S, Brown M, Ching A, Almeida R (1996) Cost-effectiveness of colorectal cancer screening in average-risk adults. In: Young G, Levin B, (eds) *Prevention and early detection of colorectal cancer*. Saunders, London, pp 321–356

37. Frazier AL, Colditz GA, Fuchs CS, Kuntz KM (2000) Cost-effectiveness of screening for colorectal cancer in the general population. *JAMA* 284:1954–1961
38. Loeve F, Brown ML, Boer R, van Ballegooijen M, van Oortmarssen GJ, Habbema JD (2000) Endoscopic colorectal cancer screening: a cost-saving analysis. *J Natl Cancer Inst* 92:557–563
39. Sonnenberg A, Delco F, Inadomi JM (2000) Cost-effectiveness of colonoscopy in screening for colorectal cancer. *Ann Intern Med* 133:373–384
40. Coffield AB, Maciosek MV, McGinnis JM, Harris JR, Caldwell MB, Teutsch SM, et al (2001) Priorities among recommended clinical preventive services. *Am J Prev Med* 21:1–9
41. Winawer S (2002) Screening for colorectal cancer. *Princ Pract Oncol* 16:1–11